Prepared for submission to JCAP

arXiv:1611.03985v1 [gr-qc] 12 Nov 2016

Homogeneous Gödel-type solutions in hybrid metric-Palatini gravity

Janilo Santos, [a,1] **Marcelo J. Rebouças,**[b] **Thiago B.R.F. Oliveira**[c] **and Antônio F.F. Teixeira**[b]

[a]Departamento de Física Teórica e Experimental
Universidade Federal do Rio Grande do Norte
59072-970 Natal – RN, Brazil

[b]Centro Brasileiro de Pesquisas Físicas
Rua Dr. Xavier Sigaud, 150
22290-180 Rio de Janeiro – RJ, Brazil

[c]Instituto Federal de Educação, Ciência e Tecnologia do Rio G. do Norte
BR 406 - km 73, 3371, Perímetro Rural
59550-000 João Câmara – RN, Brazil

[1]Corresponding author.

E-mail: janilo@dfte.ufrn.br, reboucas@cbpf.br, thiago.freitas@ifrn.edu.br, teixeira@cbpf.br

Abstract. The hybrid metric-Palatini $f(\mathcal{R})$ gravity is a recently devised approach to modified gravity in which it is added to the metric Ricci scalar R, in the Einstein-Hilbert Lagrangian, a function $f(\mathcal{R})$ of Palatini curvature scalar $\mathcal{R}$, which is constructed from an independent connection. These hybrid metric-Palatini gravity theories provide an alternative way to explain the current accelerating expansion without a dark energy matter component. If gravitation is to be described by a hybrid metric-Palatini $f(\mathcal{R})$ gravity theory there are a number of issues that ought to be examined in its context, including the question as to whether its equations allow homogeneous Gödel-type solutions, which necessarily leads to violation of causality. Here, to look further into the potentialities and difficulties of $f(\mathcal{R})$ theories, we examine whether they admit Gödel-type solutions for some physically well-motivated matter sources. We first show that under certain conditions on the matter sources the problem of finding out space-time homogeneous solutions in $f(\mathcal{R})$ theories reduces to the problem of determining solutions of this type in $f(R)$ gravity in the metric formalism. Employing this result, we determine a perfect-fluid Gödel-type solution in $f(\mathcal{R})$ gravity, and show that it is isometric to the Gödel geometry, and therefore exhibits violation causality. This extends a theorem on Gödel-type models, which was established in the framework of general relativity. We also show that a single massless scalar field gives rise to the only ST-homogeneous Gödel-type solution with no violation of causality of Gödel type. Since the perfect-fluid and scalar field solutions are in the hyperbolic family, i.e. the essential parameter is positive $m^2 > 0$, we proceed further and determine a general Gödel-type solution with a combination of a scalar with an electromagnetic field plus a perfect fluid as matter source. This general Gödel-type solution contains solutions with $m = 0$ and with $m^2 < 0$, as well as the previous solutions as special cases. The bare existence of these Gödel-type solutions makes apparent that hybrid metric-Palatini $f(\mathcal{R})$ gravity does not remedy causal anomaly in the form of closed timelike curves that are permitted in general relativity.

Keywords: gravity, modified gravity, metric-Palatini hybrid gravity, homogeneous Gödel-type solutions, violation of causality.

Contents

1 Introduction **1**

2 Hybrid metric-Palatini gravity **3**

3 Homogeneous Gödel-type geometries **4**
3.1 Homogeneity and non-equivalent metrics 5
3.2 Closed time-like curves 5

4 ST-Homogeneous solutions in hybrid metric-Palatini gravity **6**
4.1 Field equations 7
4.2 Gödel-type solutions 8
4.2.1 Perfect fluid solution 9
4.2.2 Scalar field solution 10
4.2.3 Combined-fields general solution 11

5 Concluding remarks **13**

1 Introduction

A number of cosmological observations coming from different sources, including the supernovae type Ia (SNe Ia) [1–3], the cosmic microwave background radiation (CMBR) [4, 5] and baryon acoustic oscillation (BAO) surveys [6–11], indicate that the Universe is presently expanding with an accelerating rate. The frameworks proposed to account for this observed accelerated expansion can be roughly grouped into two families. In the first, the underlying theory, general relativity (GR), is kept unchanged, and the so-called dark energy component is invoked. In this context, the simplest way to account for the accelerating expansion of the Universe is through the introduction of a cosmological constant, Λ, into Einstein's field equations. This is completely consistent with the available observational data, but it faces difficulties such as the order of magnitude of the cosmological constant and its microphysical origin. In the second family, modifications of Einstein's field equations are assumed as an alternative for explaining the accelerated expansion. This latter group includes, for example, generalized theories of gravity based upon modifications of the Einstein-Hilbert action by taking nonlinear functions, $f(R)$, of the Ricci scalar R or other curvature invariants (for reviews see Refs. [12–17]).

In dealing with $f(R)$ gravity theories two different variational approaches, which give rise to different dynamics, have often been considered in the literature [12–17]. In the so-called metric formalism the connection is assumed to be Levi-Civita, therefore defined by the metric. In the Palatini formalism the metric and the connection are treated as independent fields, and it is assumed that the matter fields do not couple with the independent connections. Although these approaches have been invoked as possible ways to satisfactorily deal with the observed accelerated expansion of the Universe, it has been pointed out that $f(R)$ gravity theories can face relevant difficulties, including the evolution of cosmological perturbations and local gravity constraints (see, for example, Refs. [15, 18–23]).

These undesirable features have motivated a recent approach to modified $f(R)$ theories of gravity, which can be employed as a possible way to handle the observed late-time cosmic acceleration, and also circumvent some difficulties that arise in the framework of $f(R)$ theories in both formalisms. The hybrid metric-Palatini $f(\mathcal{R})$ gravity is a recently devised approach to such modified theories, in which it is added to the ordinary Ricci scalar R, in the Einstein-Hilbert Lagrangian, a function, $f(\mathcal{R})$, of Palatini curvature scalar $\mathcal{R}$, which is constructed from the independent connection $\Gamma^{\rho}_{\mu\nu}$ [24]. These hybrid metric-Palatini gravity theories suitably unify the description of the late-time cosmic acceleration with the local solar system constraints [24, 25]. Some of the astrophysical and cosmological implications of hybrid metric-Palatini gravity have been examined in a number of papers [26–30]. Wormwhole solutions, Einstein static universe, linear perturbations, the Cauchy problem, dynamical system analysis and a brane model have been discussed, respectively, in the references [31–36]. Other important matters such as Noether symmetries [37] and the thermodynamic behavior [38] have also been recently considered. For an introduction to hybrid metric-Palatini $f(\mathcal{R})$ gravity and a detailed list of related references, we refer the reader to the recent review article [29].

In general relativity (GR) the space-times have *locally* the same causal structure of the flat space-time of special relativity since the space-times of GR are locally Minkowskian. On nonlocal scale, however, significant differences may arise since the general relativity field equations do no provide nonlocal constraints on the underlying space-times. Indeed, it has long been known that there are solutions to Einstein's field equations that present causal anomalies in the form of closed time-like curves (see, for example, Refs. [39–44]).

The renowned model found by Gödel [45] is the best known example of a solution to the Einstein's equations, with a physically well-motivated source, that makes it apparent that GR permits solutions with closed timelike world lines, regardless of its local Lorentzian character that ensures locally an inherited regular chronology and therefore the local validation of the causality principle. The Gödel model is a solution of Einstein's equations with cosmological constant Λ for dust of density ρ, but it can also be viewed as perfect-fluid solution with equation of state $p = \rho$ with no cosmological constant term. Owing to its unexpected properties, Gödel's solution has a recognizable importance and has motivated a considerable number of investigations on rotating Gödel-type models as well as on causal anomalies in the context of general relativity (see, e.g. Refs. [46–58]) and in the framework of other theories of gravitation (see, for example, Refs. [59–88]).

In two recent papers, we have also examined Gödel-type models and the violation of causality problem for $f(R)$ gravity in both the metric and Palatini variational approaches [75, 76], extending therefore the results of Refs. [89] and [90]. If gravitation is to be described by hybrid metric-Palatini $f(\mathcal{R})$ gravity theory there are a number of issues that ought to be reexamined in its context, including the question as to whether these gravity theories allow Gödel-type solutions, which necessarily lead to closed timelke curves, or would remedy this causal pathology by ruling out this type of solutions, which are permitted in general relativity.

In this article, to proceed further with the investigations on the potentialities, difficulties and limitations of $f(\mathcal{R})$, we undertake this question by examining whether the $f(\mathcal{R})$ gravity theories admit homogeneous Gödel-type solutions for a number of physically well-motivated matter sources. To this end, we first examine the problem of finding out ST-homogeneous solutions in hybrid metric-Palatini $f(\mathcal{R})$ gravity for matter sources with constant trace T (scalar) of the energy-momentum tensor, and show that in these cases the problem of finding

out solutions in the hybrid metric-Palatini $f(\mathcal{R})$ gravity reduces to the problem of determining ST-homogeneous solutions of $f(R)$ gravity in the metric formalism. Second, we then show that every perfect-fluid Gödel-type solution of any $f(\mathcal{R})$ gravity with density ρ and pressure p and satisfying the weak energy conditions $\rho > 0$ and $\rho + p \geq 0$ is necessarily isometric to the Gödel geometry.[1] Since any perfect-fluid ST-homogeneous Gödel-type solution of hybrid metric-Palatini gravity is noncausal and isometric to Gödel metric, the question as to whether other matter sources could generate different Gödel-type solutions naturally arises. We examine this question and show that a special single massless scalar field gives rise to the only ST-homogeneous Gödel-type solution without violation of causality of Gödel type. Since both perfect-fluid and scalar field homogeneous Gödel-type solutions of $f(\mathcal{R})$ gravity are in the hyperbolic family whose essential parameter $m^2 > 0$, the question as to whether multiple components matter sources could give rise to Gödel-type solutions with $m^2 < 0$ and with $m = 0$ naturally arises. We also examine this question by taking a combination of scalar and electromagnetic fields with a perfect fluid as a matter source, and determine a general ST-homogeneous Gödel-type solution which contains the solutions of the perfect-fluid and scalar-field solutions as particular cases, and also realizes ST-homogeneous Gödel-type solutions in the trigonometric and linear families. The bare existence of these ST-homogeneous Gödel-type solutions make apparent that hybrid metric-Palatini $f(\mathcal{R})$ gravity does not remedy causal pathology in the form of closed timelike curves that are permitted in general relativity.

The structure of the paper is as follows. In Section 2 we give a brief account of the hybrid metric-Palatini $f(\mathcal{R})$ gravity theories. In Section 3 we present the basic properties of homogenous Gödel-type geometries. This includes the ST-homogeneity conditions, a classification of ST-homogeneous Gödel-type geometries and a study of the existence of closed time-like curves in all ST-homogeneous Gödel-type metrics. In Section 4 we first examine the problem of finding out ST-homogeneous solutions in $f(\mathcal{R})$ gravity whose trace T of the energy-momentum tensor of the matter source is constant, and show that in such cases the problem of finding out solutions $f(\mathcal{R})$ gravity reduces to the problem of determining ST-homogeneous solutions of $f(R)$ gravity in the metric formalism; second, we show that the hybrid metric-Palatini $f(\mathcal{R})$ gravity theories admit ST-homogeneous Gödel-type solutions for several physically well-motivated matter contents.

2 Hybrid metric-Palatini gravity

The action that defines a hybrid metric-Palatini gravity is given by

$$S = \frac{1}{2\kappa^2}\int d^4x\sqrt{-g}\,\left[R + f(\mathcal{R}) + \mathcal{L}_m\right]\,, \tag{2.1}$$

where $\kappa^2 = 8\pi G$, g is the determinant of the metric tensor $g_{\mu\nu}$, R is the Ricci scalar associated to the Levi-Civita connection of the metric $g_{\mu\nu}$, $\mathcal{L}_m$ is the Lagragian density for the matter fields, and the extra term $f(\mathcal{R})$ is a function of Palatini curvature scalar $\mathcal{R}$, which depends on the metric and on an independent connection $\Gamma^\rho_{\mu\nu}$ through

$$\mathcal{R} \equiv g^{\mu\nu}\mathcal{R}_{\mu\nu} = g^{\mu\nu}\left(\partial_\rho\Gamma^\rho_{\mu\nu} - \partial_\nu\Gamma^\rho_{\mu\rho} + \Gamma^\rho_{\rho\lambda}\Gamma^\lambda_{\mu\nu} - \Gamma^\rho_{\mu\lambda}\Gamma^\lambda_{\rho\nu}\right). \tag{2.2}$$

[1] This extends to the context of $f(\mathcal{R})$ gravity the Bampi-Zordan theorem [91] which was established in the context of the general relativity theory, and states that every Gödel-type solution of Einstein's equations with a perfect-fluid energy-momentum tensor is necessarily isometric to the Gödel spacetime.

The variation of the action (2.1) with respect to the metric gives the field equations

$$G_{\mu\nu} + F(\mathcal{R})\mathcal{R}_{\mu\nu} - \frac{f(\mathcal{R})}{2} g_{\mu\nu} = \kappa^2 T_{\mu\nu}\,, \tag{2.3}$$

where $G_{\mu\nu} = R_{\mu\nu} - R/2\, g_{\mu\nu}$ and $R_{\mu\nu}$ are, respectively, Einstein and Ricci tensor associated with the Levi-Civita connection of $g_{\mu\nu}$, $F(\mathcal{R}) \equiv df/d\mathcal{R}$, $T_{\mu\nu} = -2/\sqrt{-g}\delta(\sqrt{-g}\mathcal{L}_m)/\delta g^{\mu\nu}$ is the energy-momentum tensor of the matter fields.

The variation of the action (2.1) with respect to the independent connection $\Gamma^\rho_{\mu\nu}$ yields

$$\widetilde{\nabla}_\beta \left(\sqrt{-g}\, F(\mathcal{R})\, g^{\mu\nu}\right) = 0\,, \tag{2.4}$$

where $\widetilde{\nabla}_\beta$ denotes the covariant derivative associated with $\Gamma^\rho_{\mu\nu}$. If one defines a metric $h_{\mu\nu} = F(\mathcal{R})\, g_{\mu\nu}$, it can be easily shown that Eq. (2.4) determines a Levi-Civita connection of $h_{\mu\nu}$, which in turn can be rewritten in terms of $g_{\mu\nu}$ and its Levi-Civita connection $\{^\rho_{\mu\nu}\}$ in the form

$$\Gamma^\rho_{\mu\nu} = \left\{^\rho_{\mu\nu}\right\} + \frac{1}{2}\left(\delta^\rho_\mu\partial_\nu + \delta^\rho_\nu\partial_\mu - g_{\mu\nu}g^{\rho\sigma}\partial_\sigma\right)\ln F(\mathcal{R})\,. \tag{2.5}$$

Using now Eq. (2.5) one finds the relation between the two Ricci tensors, which is given by

$$\mathcal{R}_{\mu\nu} = R_{\mu\nu} + \frac{3}{2F^2}\,\partial_\mu F\,\partial_\nu F - \frac{1}{F}(\nabla_\mu\nabla_\nu + \frac{1}{2}g_{\mu\nu}\Box)F\,, \tag{2.6}$$

where ∇_μ denotes the covariant derivative associated to $\{^\rho_{\mu\nu}\}$.

Equation (2.6) in turn gives rise to the following relation between the two Ricci scalars:

$$\mathcal{R} = R + \frac{3}{2F^2}(\partial F)^2 - \frac{3}{F}\Box F\,, \tag{2.7}$$

where $(\partial F)^2 = g^{\alpha\beta}\,\partial_\alpha F\partial_\beta F$ and $\Box F = g^{\alpha\beta}\,\nabla_\alpha\nabla_\beta\, F$.

The Palatini curvature $\mathcal{R}$ can be obtained from the trace of the field Eq. (2.3), which yields

$$\mathcal{R}F(\mathcal{R}) - 2f(\mathcal{R}) = \kappa^2 T + R \equiv X\,. \tag{2.8}$$

This trace equation can be used to express $\mathcal{R}$ algebraically in terms of X when the $f(\mathcal{R})$ is given as an analytic expression. Finally, we note that the variable X measures the deviation from the general relativity trace equation $R = -\kappa^2 T$.

3 Homogeneous Gödel-type geometries

To make this work clear and to a certain extent self-contained, in this section we present the basic properties of homogenous Gödel-type geometries, which we use in the following sections. To this end, we first discuss the conditions for space-time homogeneity (ST-homogeneity) of these space-times, and present all non-isometric ST-homogeneous Gödel-type classes. These ST-homogeneity conditions along with the set of isometrically non-equivalent geometries are important in the determination of ST-homogeneous Gödel-type solutions in Section 4 to the hybrid metric-Palatini $f(\mathcal{R})$ field equations. Second, we discuss the existence of closed time-like curves in the metrics of these classes. The existence of these non-causal curves are crucial to examine whether hybrid metric-Palatini $f(\mathcal{R})$ gravities allow violation of causality of Gödel-type.

3.1 Homogeneity and non-equivalent metrics

Gödel solution to the general relativity field equations is a particular member of the broad family of geometries whose general form in cylindrical coordinates, (r, ϕ, z), is given by [90]

$$ds^2 = [dt + H(r)d\phi]^2 - D^2(r)d\phi^2 - dr^2 - dz^2\,. \tag{3.1}$$

The necessary and sufficient conditions for the Gödel-type metric (3.1) to be space-time homogeneous (ST-homogeneous) are given by [90, 92]

$$\frac{H'}{D} = 2\omega \qquad \text{and} \qquad \frac{D''}{D} = m^2, \tag{3.2}$$

where the prime denote derivative with respect r, and the parameters (m, ω) are constants such that $-\infty \leq m^2 \leq \infty$ and $\omega^2 > 0$. As a matter of fact, except for the case $m^2 = 4\omega^2$ all locally ST-homogeneous Gödel-type space-times admit a group G_5 of isometries acting transitively on the whole space-time [92]. The special case $m^2 = 4\omega^2$ admits a G_7 of isometries [92, 93].

The irreducible set of isometrically nonequivalent ST-homogeneous Gödel-type metrics can be obtained by integrating equations (3.2) and suitably eliminating nonessential integration constants. The final result is that ST-homogeneous Gödel-type geometries can be grouped in the following three classes [90]:

i. Hyperbolic, in which $m^2 = \text{const} > 0$ and

$$H = \frac{4\,\omega}{m^2}\,\sinh^2(\frac{mr}{2}), \qquad D = \frac{1}{m}\sinh(mr)\,; \tag{3.3}$$

ii. Linear, in which $m = 0$ and

$$H = \omega r^2, \qquad D = r\,, \tag{3.4}$$

iii. Trigonometric, where $m^2 = \text{const} \equiv -\mu^2 < 0$ and

$$H = \frac{4\,\omega}{\mu^2}\,\sin^2(\frac{\mu r}{2}), \qquad D = \frac{1}{\mu}\sin(\mu r)\,. \tag{3.5}$$

Thus, clearly all ST-homogeneous Gödel-type geometries are characterized by the two independent parameters m^2 and ω — identical pairs (m^2, ω^2) specify isometric spacetimes [90, 92, 93].[2] In this way, to determine whether hybrid metric-Palatini $f(\mathcal{R})$ gravity allows Gödel-type solutions is to find out whether its field equations can be used to specify a pair of parameters m^2 and ω for a suitably chosen matter source.

3.2 Closed time-like curves

We begin by noting that the presence of a single closed timelike curve in a space-time is an unequivocal manifestation of violation of causality. However, a space-time may admit non-causal closed curves other than Gödel's circles we discuss in this Section. To examine the existence of closed time-like curves in ST-homogeneous Gödel-type metrics we first rewrite the line element (3.1) as

$$ds^2 = dt^2 + 2\,H(r)\,dt\,d\phi - dr^2 - G(r)\,d\phi^2 - dz^2\,, \tag{3.6}$$

[2] Gödel geometry is a solution of Eintein's equations in general relativity, is indeed a particular case of the hyperbolic class of geometries in which $m^2 = 2\omega^2$.

where $G(r) = D^2 - H^2$. In this form it is easy to show that existence of closed timelike curves, which allows for violation of causality in homogeneous Gödel-type space-times, depends on the sign of the metric function $G(r)$. Indeed, from Eq. (3.6) one has that the circles, hereafte called Gödel's circles, defined by $t, z, r = \text{const}$ become closed timelike curves whenever $G(r) < 0$.

For the hyperbolic ($m^2 > 0$) class of homogeneous Gödel-type metrics, from Eqs. (3.3) one has that

$$G(r) = \frac{4}{m^2}\,\sinh^2(\frac{mr}{2})\left[(1-\frac{4\omega^2}{m^2})\,\sinh^2(\frac{mr}{2})+1\right]. \tag{3.7}$$

Therefore for $0 < m^2 < 4\omega^2$ there is a critical radius r_c defined by $G(r) = 0$, which is given by

$$\sinh^2\frac{mr_c}{2} = \left[\frac{4\omega^2}{m^2}-1\right]^{-1}, \tag{3.8}$$

such that for $r < r_c$ one has $G(r) > 0$, and for $r > r_c$ one has $G(r) < 0$. Thus, the circles $t, r, z = \text{const}$ with $r > r_c$ are closed timelike curves.[3]

For linear class ($m = 0$) of homogeneous Gödel-type space-times, from Eq. (3.4) one easily finds

$$G(r) = r^2 - r^4\,\omega^2 = -r^2\,(r\,\omega - 1)\,(r\,\omega + 1)\,. \tag{3.9}$$

Thus, there is a critical radius, defined by $G(r) = 0$, and given by $r_c = 1/\omega$, such that for any radius $r > r_c$ one has $G(r) < 0$, and then the circles defined by $t, z, r = \text{const}$ are closed timelike curves.

Finally for the trigonometric class ($m^2 = \text{const} \equiv -\mu^2 < 0$), from the metric functions given by Eq. (3.5) one finds

$$G\,(r) = \frac{4}{\mu^4}\,\sin^2(\frac{\mu\,r}{2})\,[\,\mu^2 - (4\,\omega^2+\mu^2)\,\sin^2(\frac{\mu\,r}{2})\,]\,, \tag{3.10}$$

and therefore $G(r)$ has an infinite sequence of zeros. Thus, in the section $t, z, r = \text{const}$, there is an sequence of alternating causal $[\,G(r) > 0\,]$ and noncausal $[\,G(r) < 0\,]$ regions without and with noncausal circles, depending on the value of $r = \text{const}$ (For more details see the Appendix of Ref. [82]). In this way, if $G(r) < 0$ for a certain range of r ($r_1 < r < r_2$, say) noncausal Gödel's circles exist, whereas for r in the next circular band $r_2 < r < r_3$ (say) for which $G(r) > 0$ no such closed timelike circles exist, and so on.

To close this section, we note that in this paper by non-causal and causal solutions we mean, respectively, solutions with and without violation of causality of Gödel-type, i.e., with and without Gödel's circles.

4 ST-Homogeneous solutions in hybrid metric-Palatini gravity

The aim of this section is twofold. First, we examine the problem of finding out ST-homogeneous solutions in hybrid metric-Palatini $f(\mathcal{R})$ gravity whose trace T of the energy-momentum tensor of the matter source is constant. We show that in such cases the problem of finding out solutions in the hybrid metric-Palatini $f(\mathcal{R})$ gravity reduces to the problem of determining ST-homogeneous solutions of $f(R)$ gravity in the metric formalism. Second, we

[3]The only ST-homogeneous Gödel-type space-time without these noncausal circles come about when $m^2 = 4\omega^2$ (see Ref. [90]). In this case, the critical radius $r_c \to \infty$, and hence the violation of causality of Gödel type is avoided.

examine whether hybrid metric-Palatini $f(\mathcal{R})$ field equations admit ST-homogeneous Gödel-type solutions for several physically well-motivated matter contents.

4.1 Field equations

We begin by noting that using equation (2.6) the field equations (2.3) of the hybrid metric-Palatini $f(\mathcal{R})$ gravity can be rewritten in the form

$$G_{\mu\nu} = \kappa^2 \, T^{\text{eff}}_{\mu\nu} = \kappa^2 \left(T_{\mu\nu} + T^{\mathcal{R}}_{\mu\nu} \right) , \tag{4.1}$$

where

$$\kappa^2 T^{\mathcal{R}}_{\mu\nu} = \frac{1}{2} \left[f(\mathcal{R}) + \Box F(\mathcal{R}) \right] g_{\mu\nu} - F(\mathcal{R}) R_{\mu\nu} + \nabla_\mu \nabla_\nu F(\mathcal{R}) - \frac{3}{2F(\mathcal{R})} \, \partial_\mu F(\mathcal{R}) \partial_\nu F(\mathcal{R}). \tag{4.2}$$

In this context, an important constraint comes from the trace of the field equations (4.1), which can be written in the form

$$R + \kappa^2 T = X = -\kappa^2 T^{\mathcal{R}} , \tag{4.3}$$

where from equation (4.2) one has

$$\kappa^2 T^{\mathcal{R}} = 2f(\mathcal{R}) - RF(\mathcal{R}) + 3\Box F(\mathcal{R}) - \frac{3}{2} \frac{[\partial F(\mathcal{R})]^2}{F(\mathcal{R})} . \tag{4.4}$$

For ST-homogeneous spacetimes, which we are concerned with in this paper, one has that the Ricci scalar is necessarily constant. On the other hand, for matter sources whose trace of the energy-momentum T is also constant, which we focus in this paper, one has $X = \text{const} = -\kappa^2 T^{\mathcal{R}}$. In such cases, one can easily show that the field equations of the hybrid metric-Palatini gravity reduce to field equations of $f(R)$ theories in the metric formalism.

To this end, we first examine the second term on the right hand side of equation (2.5) which gives the departures of the independent connection $\Gamma^\rho_{\mu\nu}$ from Levi-Civita connection $\{^\rho_{\mu\nu}\}$. Clearly, each individual part of this second term in this equation is proportional to

$$\partial_\mu \ln F(\mathcal{R}) = \frac{F'}{F} \, \partial_\mu \mathcal{R} = \frac{1}{F} \, \partial_\mu F(\mathcal{R}) . \tag{4.5}$$

On the other hand, to calculate $\partial_\mu F(\mathcal{R})$ we note that from the trace equation (2.8) one has $\partial_\mu \mathcal{R} = \partial_\mu X \, / \, [\mathcal{R}F'(\mathcal{R}) - F(\mathcal{R})]$, which together with equation (4.5) furnishes

$$\partial_\mu F(\mathcal{R}) = \frac{F'(\mathcal{R}) \, \partial_\mu X}{\mathcal{R}F'(\mathcal{R}) - F(\mathcal{R})} , \tag{4.6}$$

provided that $\mathcal{R}F'(\mathcal{R}) - F(\mathcal{R}) \neq 0$.[4] From equations (4.5) and (4.6) one has that for $X = \text{const}$ the connection $\Gamma^\rho_{\mu\nu}$ reduces to Levi-Civita connection $\{^\rho_{\mu\nu}\}$. Furthemore, from equations (2.6) and (2.7) one can easily show that $\partial_\mu F(\mathcal{R}) = 0$ also ensures that $\mathcal{R}_{\mu\nu} = R_{\mu\nu}$ and $\mathcal{R} = R$. This makes apparent that ST-homogeneous spacetimes solutions whose trace of the energy-momentum tensor is constant ($R + \kappa^2 T = X = \text{const}$) the field equations of the

[4] This inequality holds as long as $f(\mathcal{R})$ is a homogeneous function of degree $n \neq 2$. Particularly for $n = 2$, by virtue of Euler's theorem for homogeneous functions, the left hand side of the Eq. (2.8) vanishes, thus $X = 0$ regardless of the pair $(\mathcal{R}, T)$. In this case ($n = 2$) the function $f(\mathcal{R})$ is said to be degenerate. For any other functional form for $f(\mathcal{R})$ this inequality ought to be examined.

hybrid metric-Palatini gravity (2.3) reduce to field equations of $f(R)$ theories in the metric formalism, which can clearly be rewritten in the form

$$[1+F(R)]\,G_{\mu\nu}-\frac{1}{2}\,[\,f(R)-RF(R)\,]\,g_{\mu\nu}=\kappa^2\,T_{\mu\nu}\,, \tag{4.7}$$

with associated trace equation

$$RF(R)-2f(R)=\kappa^2 T+R=\text{const.} \tag{4.8}$$

4.2 Gödel-type solutions

In this section we discuss ST-homogeneous Gödel-type solutions in hybrid metric-Palatini $f(\mathcal{R})$ gravity for well-motivated matter contents whose trace of the energy-momentum tensor is constant.

We begin by noting that the search for ST-homogeneous Gödel-type solutions to the hybrid metric-Palatini gravity field equations is greatly simplified if instead of using coordinates basis one uses a new basis given by the following set of linearly independent one-forms (tetrad frame) Θ^A:

$$\theta^0=dt+H(r)d\phi\,,\quad \theta^1=dr\,,\quad \theta^2=D(r)d\phi\,,\quad \theta^3=dz\,, \tag{4.9}$$

relative to which the Gödel-type line element (3.1) takes the local Lorentzian form

$$ds^2=\eta_{AB}\,\theta^A\theta^B=(\theta^0)^2-(\theta^1)^2-(\theta^2)^2-(\theta^3)^2\,. \tag{4.10}$$

Here and in what follows capital letters are tetrad indices (or Lorentz frame indices) and run from 0 to 3. These Lorentz frame indices are raised and lowered with Lorentz matrices $\eta^{AB}=\eta_{AB}=\text{diag}(+1,-1,-1,-1)$, respectively.

In the tetrad frame (4.9) the nonvanishing components of the Ricci tensor, $R_{AB}=\eta^{CD}R_{CADB}$, are given by

$$R_{02}=\frac{1}{2}\left(\frac{H'}{D}\right)',\qquad R_{00}=\frac{1}{2}\left(\frac{H'}{D}\right)^2, \tag{4.11}$$

$$R_{11}=R_{22}=\frac{1}{2}\left(\frac{H'}{D}\right)^2-\frac{D''}{D}\,, \tag{4.12}$$

where the prime denotes derivative with respect to r. Since the Lorentz frame components of the Ricci tensor depend only on H'/D and D''/D, from the ST-homogeneity conditions (3.2) one has that for all classes of ST-homogeneous Gödel-type metrics the frame components of the Ricci tensor are constants. Thus, the Ricci scalar is also constant and given by $R=\eta^{CD}R_{CD}=2(m^2-\omega^2)$.

In this Lorentzian basis the field equations (4.7) reduce to

$$[1+F(R)]\,G_{AB}-\frac{1}{2}\,[f(R)-RF(R)]\,\eta_{AB}=\kappa^2\,T_{AB}\,, \tag{4.13}$$

where from equations (4.11) and (4.12) along with the conditions (3.2) one has that the only nonvanishing Lorentz frame components of the Einstein tensor G_{AB} for ST-homogeneous Gödel-type metrics take the very simple form

$$G_{00}=3\omega^2-m^2\,,\qquad G_{11}=G_{22}=\omega^2\,,\qquad G_{33}=m^2-\omega^2\,. \tag{4.14}$$

Before proceeding to the exam of ST-homogeneous Gödel-type solutions in hybrid metric-Palatini gravity for different matter contents, some words of clarification concerning previous results in the literature are in order here. Since the problem of finding out ST-homogeneous solutions in hybrid metric-Palatini $f(\mathcal{R})$ gravity for matter source with trace $T = \text{const}$ reduces to the problem of determining ST-homogeneous solutions of $f(R)$ gravity in the metric formalism, the perfect fluid and scalar field Gödel-type solutions to the $f(R)$ gravity field equations found in Ref. [75] are clearly solutions in hybrid metric-Palatini $f(\mathcal{R})$ gravity. In the following, besides present new ST-homogeneous Gödel-type solutions, we shall also recover some of these known solutions but using the notation of the present work for subsequent comparison with new results.

4.2.1 Perfect fluid solution

In the Lorentzian basis (4.9) the energy-momentum tensor of a perfect fluid of density ρ and pressure p is given by

$$T^{(M)}_{AB} = (\rho + p)\, u_A u_B - p\, \eta_{AB}\,. \tag{4.15}$$

Thus, taking into account (4.14), for this matter source one can easily show that the field equations (4.13) reduces to

$$\kappa^2\rho = 2\omega^2 F(R) + 3\omega^2 - m^2 - \frac{1}{2}f(R)\,, \tag{4.16}$$

$$\kappa^2 p = (2\omega^2 - m^2)F(R) + \omega^2 + \frac{1}{2}f(R)\,, \tag{4.17}$$

$$\kappa^2 p = \frac{1}{2}f(R) + m^2 - \omega^2\,. \tag{4.18}$$

The compatibility of Eq. (4.17) and Eq (4.18) leads to

$$(2\omega^2 - m^2)\,[1 + F(R)] = 0\,. \tag{4.19}$$

To ensure that the effective Newton constant $G_{\text{eff}} = G/[1 + F(R)]$ does not change its sign (see, e.g., Refs. [19, 94] for details) we assume that $1 + F(R) > 0$. Therefore, Eq. (4.19) gives $m^2 = 2\omega^2$, which defines the Gödel metric, and the remaining field equations, Eqs. (4.16) and (4.17), reduce to

$$\kappa^2\rho = m^2 F(R) - \frac{1}{2}f(R) + \frac{1}{2}m^2\,, \tag{4.20}$$

$$\kappa^2 p = \frac{1}{2}f(R) + \frac{1}{2}m^2\,, \tag{4.21}$$

which furnish

$$m^2 = 2\omega^2 = \frac{\kappa^2(\rho + p)}{1 + F(R)}\,, \tag{4.22}$$

making explicit how the $f(R)$ gravity can be used to specify a pair of parameters (m^2, ω^2), thus a ST-homogeneous Gödel-type solution, for a perfect fluid of density ρ and pressure p, which satisfy the weak energy conditions $\rho > 0$ and $\rho + p \geq 0$.

Finally, regarding the causality properties of these solutions, since they are isometric to Gödel geometry, they inevitably exhibit closed time-like Gödel's circles. The critical radius r_c is given by eq. (3.8), which taking into account equation (4.22) can be rewritten as

$$r_c = \frac{2}{m}\,\sinh^{-1}(1) = 2\,\sinh^{-1}(1)\,\sqrt{\frac{1 + F(R)}{\kappa^2(\rho + p)}}\,, \tag{4.23}$$

making apparent that the critical radius, beyond which noncausal Gödel's circles exist, depends on the gravity theory and the matter content as one would have expected from the outset.

4.2.2 Scalar field solution

Since any perfect-fluid ST-homogeneous Gödel-type solution of hybrid metric-Palatini gravity is noncausal and isometric to Gödel metric, the question as to whether other matter sources could generate different Gödel-type solutions naturally arises here. In this section we shall examine whether a single scalar field can be matter source of a ST-homogeneous Gödel-type solution in hybrid metric-Palatini $f(\mathcal{R})$ gravity.

In the Lorentzian basis (4.9) the energy-momentum tensor of a single scalar field is given by

$$T^{(S)}_{AB} = \Phi_{|A}\,\Phi_{|B} - \frac{1}{2}\,\eta_{AB}\,\Phi_{|M}\,\Phi_{|N}\,\eta^{MN}, \tag{4.24}$$

where a vertical bar denotes components of covariant derivatives relative to the local basis $\theta^A = e^A_{\ \alpha}\,dx^\alpha$ [see Eqs. (4.9) and (4.10)], i.e. $\Phi_{|A} = e^A_{\ \mu}\,\nabla_\mu\Phi$. Following Ref. [90] it is straightforward to show that a scalar field of the form $\Phi(z) = \varepsilon\,z + \epsilon$, with $\varepsilon, \epsilon = \text{const}$, fulfills the scalar field equation $\Box\,\Phi = \eta^{AB}\,\nabla_A\nabla_B\,\Phi = 0$. Thus, the non-vanishing components of the energy-moment tensor for this scalar field are

$$T^{(S)}_{00} = -T^{(S)}_{11} = -T^{(S)}_{22} = T^{(S)}_{33} = \frac{\varepsilon^2}{2}\,, \tag{4.25}$$

and the field equations (4.13) can be written in the form

$$k^2\varepsilon^2 = 4\omega^2 F(R) - f(R) + 2(3\omega^2 - m^2)\,, \tag{4.26}$$

$$k^2\varepsilon^2 = 2(m^2 - 2\omega^2)F(R) - f(R) - 2\omega^2\,, \tag{4.27}$$

$$k^2\varepsilon^2 = f(R) + 2(m^2 - \omega^2)\,. \tag{4.28}$$

The compatibility of Eqs. (4.26) and (4.27) implies

$$(m^2 - 4\omega^2)[1 + F(R)] = 0\,. \tag{4.29}$$

Assuming $1 + F(R) > 0$ to keep the sign of G_{eff} [19, 94], from Eq. (4.29) one has $m^2 = 4\omega^2$, which defines the ST-homogeneous hyperbolic Gödel-type geometry (3.3) with no closed Gödel circles since $r_c \to \infty$ [see Eq. (3.8)]. The remaining field equations, Eqs. (4.26) and (4.28), reduce to

$$k^2\varepsilon^2 = m^2 F(R) - f(R) - \frac{m^2}{2}\,, \tag{4.30}$$

$$k^2\varepsilon^2 = f(R) + \frac{3m^2}{2}\,, \tag{4.31}$$

which furnish

$$m^2 = 4\omega^2 = \frac{2\,\kappa^2\varepsilon^2}{1 + F(R)}\,, \tag{4.32}$$

making explicit how the $f(R)$ gravity can be used to specify a pair of parameters (m^2, ω^2), thus the ST-homogeneous Gödel-type solution for a scalar field source $\Phi(z) = \varepsilon\,z + \epsilon$, with no Gödel circles.

4.2.3 Combined-fields general solution

Since both perfect-fluid and scalar field homogeneous Gödel-type solutions of hybrid metric-Palatini $f(\mathcal{R})$ gravity are in the hyperbolic family $m^2 > 0$ [cf. Eq.(3.3)], the question as to whether multiple components matter sources could give rise to Gödel-type solutions in the trigonometric [$m^2 = -\mu^2 < 0$, cf. Eq. (3.5)] and linear [$m = 0$, cf. Eq. 3.4)] classes naturally arises here. In this section we examine this question. To this end, we take combination of scalar and electromagnetic fields with a perfect fluid as a matter source, and find a general ST-homogeneous Gödel-type solution, which contains the solutions of the previous Sec. 4.2.1 and Sec. 4.2.2 as particular cases, and also realizes ST-homogeneous Gödel-type solutions in the trigonometric and linear families.

In the Lorentzian basis (4.9) the energy-momentum tensor of combined matter sources takes the form

$$T_{AB} = T^{(M)}_{AB} + T^{(S)}_{AB} + T^{(EM)}_{AB}\,, \tag{4.33}$$

where the non-vanishing components of $T^{(M)}_{AB}$ and $T^{(S)}_{AB}$ are given, respectively, by equations (4.15) and (4.24). Following Ref. [90], the electromagnetic field tensor F_{AB} given by

$$F_{03} = -F_{30} = E_0 \sin[2\omega(z - z_0)]\,, \qquad F_{12} = -F_{21} = E_0 \cos[2\omega(z - z_0)]\,, \tag{4.34}$$

satisfies the source-free Maxwell equations which, in the tetrad frame (4.9), take the form

$$F^{AB}{}_{|B} + \gamma^{A}{}_{MB}F^{MB} + \gamma^{C}{}_{MC}F^{AM} = 0, \qquad F_{[AB|C]} + 2F_{M[C}\gamma^{M}{}_{AB]} = 0\,, \tag{4.35}$$

where the brackets denote total anti-symmetrization and the Ricci rotation coefficients are defined by $\gamma^{A}{}_{BC} = -\nabla_\beta \theta^A_\alpha\, \theta^\alpha_B\, \theta^\beta_C$. The non-vanishing components of the associated energy-momentum tensor $T^{(EM)}_{AB} = F_A{}^C\, F_{BC} + \frac{1}{4}\eta_{AB}F^{CD}F_{CD}$ are given by

$$T^{(EM)}_{00} = T^{(EM)}_{11} = T^{(EM)}_{22} = -T^{(EM)}_{33} = \frac{E_0^2}{2}\,. \tag{4.36}$$

Thus, taking into account equations (4.14), (4.15), (4.25) and (4.36), one has that for the combined-fields matter source (4.33) the field equations (4.13) reduce to

$$2\omega^2 F(R) - \frac{1}{2}\,f(R) + 3\omega^2 - m^2 = \kappa^2\left(\rho + \frac{\varepsilon^2}{2} + \frac{E_0^2}{2}\right)\,, \tag{4.37}$$

$$(2\omega^2 - m^2)F(R) + \frac{1}{2}\,f(R) + \omega^2 = \kappa^2\left(p - \frac{\varepsilon^2}{2} + \frac{E_0^2}{2}\right)\,, \tag{4.38}$$

$$\frac{1}{2}\,f(R) + m^2 - \omega^2 = \kappa^2\left(p + \frac{\varepsilon^2}{2} - \frac{E_0^2}{2}\right)\,. \tag{4.39}$$

From Eq. (4.39) one has

$$f(R) = 2\omega^2 - 2m^2 + \kappa^2(2p + \varepsilon^2 - E_0^2)\,. \tag{4.40}$$

Substituting (4.40) into (4.37) one finds

$$2\omega^2 = \frac{\kappa^2(\rho + p + \varepsilon^2)}{1 + F(R)}\,, \tag{4.41}$$

where here and in what follows we again assume that $1 + F(R) > 0$, which is the condition to keep the sign of $G_{\rm eff}$ [19, 94] in $f(R)$ gravity theories.

To determine the other essential parameter m^2, we substitute (4.40) into (4.38) and use (4.41) to find

$$m^2 = \frac{\kappa^2(\rho + p + 2\varepsilon^2 - E_0^2)}{1 + F(R)}\,. \tag{4.42}$$

Since ST-homogeneous Gödel-type geometries are characterized by the two essential parameters m^2 and ω^2, the above equations (4.41) and (4.42) make explicit how the $f(\mathcal{R})$ gravity specifies a pair of parameters (m^2, ω^2), and therefore determines a general ST-homogeneous Gödel-type solution, for the combined-fields matter source (4.33).

The general solution given by equations (4.41) and (4.42) contains the previous solutions (4.22) and (4.32) as particular cases. Indeed, the perfect fluid Gödel solution (4.22) is recovered when $\varepsilon = E_0 = 0$ with $\rho, p \neq 0$, whereas the scalar field causal solution (4.32) is retrieved for $\rho = p = E_0 = 0$ with $\varepsilon \neq 0$.[5]

Finally, from equation (4.42) one has that the combination of scalar plus electromagnetic fields with a perfect fluid gives rise to ST-homogeneous Gödel-type solutions in hyperbolic class ($m^2 > 0$) when $E_0^2 \leq \rho + p + 2\varepsilon^2$, in the linear family ($m^2 = 0$) for $E_0^2 = \rho + p + 2\varepsilon^2$, and also in the trigonometric class ($m^2 < 0$) when $E_0^2 > \rho + p + 2\varepsilon^2$. Moreover, from equation (4.42) we also have that these three families of ST-homogeneous Gödel-type solutions can also be generated with a simple combination of scalar and electromagnetic fields (thus for $\rho = p = 0$), depending on the relative values of the amplitudes ε^2 and E_0^2, which give the sign of the term $2\varepsilon^2 - E_0^2$ in equation (4.42).

As a concrete example, we consider the $f(\mathcal{R})$ theory

$$f(\mathcal{R}) = \frac{\alpha}{\mathcal{R}^n}\,, \tag{4.43}$$

where α and n are free parameters to be determined by local gravity constraints and cosmological observations. For this $f(\mathcal{R})$ gravity, the equations (4.41) and (4.42), that specify the parameters m^2 and ω^2 for the combined-fields matter source (4.33), can be rewritten, respectively, as

$$2\omega^2[2(m^2 - \omega^2)]^{n+1} - \kappa^2(\rho + p + \varepsilon^2)\,[2(m^2 - \omega^2)]^{n+1} - 2n\alpha\omega^2 = 0\,, \tag{4.44}$$

$$m^2[2(m^2 - \omega^2)]^{n+1} - \kappa^2(\rho + p + 2\varepsilon^2 - E_0^2)\,[2(m^2 - \omega^2)]^{n+1} - n\alpha m^2 = 0\,, \tag{4.45}$$

where we have used that $F(\mathcal{R}) = df/d\mathcal{R}$, and that $\mathcal{R} = R = 2(m^2 - \omega^2)$ for these families of ST-homogeneous Gödel-type solutions. Thus, for example, to determine m^2 for the Gödel solution ($m^2 = 2\omega^2$) generated by the combination of a perfect fluid with a scalar plus an electromagnetic fields ($\varepsilon = E_0 \neq 0$, see footnote 5), from equations (4.44) and (4.45) one has to find the positive roots m^2 of the algebraic equation

$$m^{2n+2} - \kappa^2(\rho + p + \varepsilon^2)\,m^{2n} - n\alpha = 0\,. \tag{4.46}$$

Similarly, to find out m^2 for the causal ST-homogeneous Gödel solution $m^2 = 4\omega^2$, generated by a single scalar field matter source ($\rho = p = E_0 = 0$, $\varepsilon \neq 0$), we have to find the positive roots $m^2 > 0$ of the algebraic equation

$$m^{2n+2} - 2\kappa^2\varepsilon^2\,m^{2n} - \left(\frac{2}{3}\right)^{n+1} n\alpha = 0\,. \tag{4.47}$$

[5] We note that Gödel metric ($m^2 = 2\omega^2$) can also be generated through a particular combination of perfect fluid with a scalar field, namely for non-vanishing equal amplitude of the scalar and electromagnetic fields $\varepsilon = E_0 \neq 0$. Indeed in this case equations (4.41) and (4.42) give $m^2 = 2\omega^2 = [\kappa^2(\rho + p + \varepsilon^2)]/[1 + F(R)]$.

5 Concluding remarks

Despite the great success of the general relativity theory, a good deal of effort has recently gone into the study of the so-called modified gravity theories. In the cosmological modeling framework, this is motivated by the fact that these theories provide an alternative way to explain the current accelerating expansion of the Universe with no need to invoke the dark energy matter component. The hybrid metric-Palatini $f(\mathcal{R})$ gravity is a recently devised approach to such modified theories, in which it is added to the ordinary Ricci scalar R, in the Einstein-Hilbert Lagrangian, a function, $f(\mathcal{R})$, of Palatini curvature scalar $\mathcal{R}$, which is constructed from the independent connection $\Gamma^{\rho}_{\mu\nu}$.

In general relativity, the bare existence of the Gödel solution to Einstein's equations, for a physically well-motivated perfect-fluid source, makes it apparent that this theory permits solutions with violation of causality on nonlocal scale, regardless of its local Lorentzian character that ensures the local validation of the causality principle. In the context of the hybrid metric-Palatini $f(\mathcal{R})$ gravity theories the space-time manifolds are also assumed to be locally Lorentzian. Hence the chronology and causality structure of special relativity are inherited locally. The nonlocal question, however, is left open, and violation of causality can in principle arise.

Since homogeneous Gödel-type geometries necessarily lead to the existence of closed timelike circles (Section 3.2), which is an unequivocal manifestation of violation of causality, a natural way to tackle this question is by investigating whether the hybrid metric-Palatini $f(\mathcal{R})$ gravity theories permit Gödel-type solutions to their field equations. Furthermore, if gravity is to be governed by a $f(\mathcal{R})$ there are a number of issues ought to be reexamined in its context, including the question as to whether these theories admit Gödel-type solutions, or would remedy the violation of causality problem by ruling out this type of solutions, which are permitted in general relativity.

In this article, to proceed further with the investigations on the potentialities, difficulties and limitations of $f(\mathcal{R})$, we have examined whether $f(\mathcal{R})$ gravity theories admit homogeneous Gödel-type solutions for some physically well-motivated matter sources. In this way, we have first examined the problem of finding out ST-homogeneous solutions in these hybrid gravity theories whose trace T (invariant) of the energy-momentum tensor is constant. We have shown that under this assumption the problem of finding out solutions in the hybrid metric-Palatini $f(\mathcal{R})$ gravity reduces to the problem of determining ST-homogeneous solutions of $f(R)$ gravity in the metric formalism. This simplifies considerably the determination of ST-homogeneous solutions in $f(\mathcal{R})$. Employing this simplifying result, we first have found perfect-fluid Gödel-type solution of $f(\mathcal{R})$ gravity, which is necessarily isometric to the Gödel geometry, and therefore exhibit violation of causality. This extends to the context of $f(\mathcal{R})$ gravity the Bampi-Zordan theorem [91] which was established in the context of the general relativity theory.

As any perfect-fluid ST-homogeneous Gödel-type solution of hybrid metric-Palatini gravity is isometric to Gödel metric and hence noncausal, the question as to whether other matter sources could generate inequivalent solutions naturally arises. We have examined this question and shown that a single massless scalar field gives rise to the only ST-homogeneous Gödel-type solution without violation of causality of Gödel type.

Since both perfect-fluid and scalar field homogeneous Gödel-type solutions of $f(\mathcal{R})$ gravity are in the hyperbolic family whose essential parameter of Gödel-type geometries is positive ($m^2 > 0$), we have also examined whether a combination of scalar and electromagnetic fields

with a perfect fluid could be matter source of Gödel-type solutions with $m^2 < 0$ (trigonometric family) and with $m = 0$ (linear class). A general ST-homogeneous Gödel-type solution, which contains the solutions of the perfect-fluid and scalar-field solutions as particular cases, has hence been found. We have also shown that particular cases of this general solution are ST-homogeneous Gödel-type solutions in the trigonometric ($m^2 < 0$) and linear ($m = 0$) families.

To conclude, we emphasize that the bare existence of these ST-homogeneous Gödel-type solutions that we have found makes apparent that the hybrid metric-Palatini $f(\mathcal{R})$ gravity does not remedy causal pathology in the form of closed timelike curves that arises in the context of general relativity. In this regard, we finally recall the chronology protection conjecture introduced by Hawking [95], which suggests that even though closed timelike curves are classically possible, quantum effects are likely to prevent such time travel. In this way, a proper quantum gravity theory would prevent closed timelike curves from appearing.

Acknowledgments

M.J. Rebouças acknowledges the support of FAPERJ under a CNE E-26/102.328/2013 grant and also thanks CNPq for the grant under which this work was carried out. J. Santos acknowledges support of Programa de Pós-Graduação em Física - CCET/UFRN.

References

[1] A.G. Riess et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, *Astron. J.* **116** (1998) 1009.

[2] S. Perlmutter, et al., *Discovery of a supernova explosion at half the age of the Universe*, *Nature* **391** (1998) 51.

[3] S. Perlmutter et al., *Measurements of* Ω *and* Λ *from 42 high-redshift supernovae*, *Astrophys. J.* **517** (1999) 565.

[4] R. Adam et al., *Planck 2015 results. I. Overview of products and scientific results*, *Astron. & Astrophys.* **594** (2016) A1.

[5] D.N. Spergel et al., *Three-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Implications for Cosmology*, *Astrophys. J. Suppl.* **170** (2007) 377S.

[6] S. Cole et al., *The 2dF Galaxy Redshift Survey: Power-spectrum analysis of the final data set and cosmological implications*, *Mon. Not. Roy. Astron. Soc.* **362** (2005) 505.

[7] D.J. Eisenstein et al., *Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies*, *Astrophys. J.* **633** (2005) 560.

[8] W.J. Percival et al., *Baryon acoustic oscillations in the Sloan Digital Sky Survey Data Release 7 galaxy sample*, *Mon. Not. Roy. Astron. Soc.* **401** (2010) 2148.

[9] N. Padmanabhan et al., *A 2 per cent distance to* $z = 0.35$ *by reconstructing baryon acoustic oscillations - I. Methods and application to the Sloan Digital Sky Survey*, *Mon. Not. Roy. Astron. Soc.* **427** (2012) 2132.

[10] C. Blake et al, *The Wiggle Z Dark Energy Survey: mapping the distance-redshift relation with baryon acoustic oscillations*, *Mon. Not. Roy. Astron. Soc.* **418** (2011) 1707.

[11] L. Anderson et al, *The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations in the Data Release 9 Spectroscopic Galaxy Sample*, *Mon. Not. Roy. Astron. Soc.* **428** (2013) 1036.

[12] A. De Felice and S. Tsujikawa, *f(R) Theories*, *Living Rev. Rel.* **13** (2010) 3.

[13] T.P. Sotiriou and V. Faraoni, *f(R) theories of gravity*, *Rev. Mod. Phys.* **82** (2010) 451.

[14] S. Nojiri and S.D. Odintsov, *Unified cosmic history in modified gravity: From F(R) theory to Lorentz non-invariant models*, *Phys. Rep.* **505** (2011) 59.

[15] G.J. Olmo, *Palatini approach to modified gravity: f(R) theories and beyond*, *Int. J. Mod. Phys. D*, **20** (2011) 413.

[16] S. Capozziello and M. De Laurentis, *Extended Theories of Gravity*, *Phys. Rep.* **509** (2011) 167.

[17] S. Capozziello and V. Faraoni, *Beyond Einstein Gravity*, *Fundamental Theories of Physics* **170**, (Springer, Dordrecht, 2011).

[18] W. Hu and I. Sawicki, *Models of $f(R)$ cosmic acceleration that evade solar system tests*, *Phys. Rev. D* **76** (2007) 064004.

[19] S. Tsujikawa, *Observational signatures of $f(R)$ dark energy model that satisfy cosmological and local gravity constraints*, *Phys. Rev. D* **77** (2008) 023507.

[20] T. Koivisto and H. Kurki-Suonio, *Cosmological perturbations in the Palatini formulation of modified gravity*, *Class. Quantum Grav.* **23** (2006) 2355.

[21] T. Koivisto, *The matter power spectrum in $f(R)$ gravity*, *Phys. Rev. D* **73** (2006) 083517.

[22] G.J. Olmo, *Violation of the equivalence principle in modified theories of gravity*, *Phys. Rev. Lett.* **98** (2007) 061101.

[23] G.J. Olmo, *Hydrogen atom in Palatini theories of gravity*, *Phys. Rev. D* **77** (2008) 084021.

[24] T. Harko, T.S. Koivisto, F.S.N. Lobo and G.J. Olmo, *Metric-Palatini gravity unifying local constraints and late-time cosmic acceleration*, *Phys. Rev. D* **85** (2012) 084016.

[25] S. Capozziello, T. Harko, F.S.N. Lobo and G.J. Olmo, *Hybrid modified gravity unifying local tests, galactic dynamics and late-time cosmic acceleration*, *Int. J. Modern Phys. D* **22** (2013) 1342006.

[26] S. Capozziello et al., *Cosmology of hybrid metric-Palatini f(X)-gravity*, *J. Cosmol. Astropart. Phys.* **04** (2013) 011.

[27] S. Capozziello et al., *The virial theorem and the dark matter problem in hybrid metric-Palatini gravity*, *J. Cosmol. Astropart. Phys.* **07** (2013) 024.

[28] S. Capozziello et al., *Galactic rotation curves in hybrid metric-Palatini gravity*, *Astropart. Phys.* **50** (2013) 65.

[29] S. Capozziello et al., *Hybrid Metric-Palatini Gravity*, *Universe*, **1** (2015) 199.

[30] N.A. Lima and V.S. Barreto, *Constraints on hybrid metric-Palatini gravity from background evolution*, *Astrophys. J.* **818** (2016) 186.

[31] S. Capozziello et al., *Wormholes supported by hybrid metric-Palatini gravity*, *Phys. Rev. D* **86** (2012) 127504.

[32] C.G. Böhmer, F.S.N. Lobo and N. Tamanini, *Einstein static universe in hybrid metric-Palatini gravity*, *Phys. Rev. D* **88** (2013) 104019.

[33] N.A. Lima, *Dynamics of linear perturbations in the hybrid metric-Palatini gravity*, *Phys. Rev. D* **89** (2014) 083527.

[34] S. Capozziello et al., *The Cauchy problem in hybrid metric-Palatini $f(X)$-gravity*, *Int. J. Geom. Methods Mod. Phys.* **11** (2014) 1450042.

[35] S. Carloni, T. Koivisto and F.S.N. Lobo, *Dynamical system analysis of hybrid metric-Palatini cosmologies*, *Phys. Rev. D* **92** (2015) 064035.

[36] Qi-Ming Fu et al., *Hybrid metric-Palatini brane system*, *Phys. Rev. D* **94** (2016) 024020.

[37] A. Borowiec et al., *Invariant solutions and Noether symmetries in hybrid gravity*, *Phys. Rev. D* **91** (2015) 023517.

[38] T. Azizi and N. Borhani, *Thermodynamics in hybrid metric-Palatini gravity*, *Astrophys. Space Sci.* **357** (2015) 146.

[39] W.J. van Stockum, *The gravitational field of a distribution of particles rotating around an axis of symmetry*, *Proc. R. Soc. Edin.* **57** (1937) 135.

[40] M.S. Morris, K.S. Thorne and U. Yurtsever, *Wormholes, time machines, and the weak energy condition*, *Phys. Rev. Lett.* **61** (1988) 1446.

[41] M.S. Morris and K.S. Thorne, *Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity*, *Am. J. Phys.* **56** (1988) 395.

[42] F.J. Tipler, *Rotating cylinders and the possibility of global causality violation*, *Phys. Rev. D* **9** (1974) 2203.

[43] J.R. Gott III, *Closed timelike curves produced by pairs of moving cosmic strings: Exact solutions*, *Phys. Rev. Lett.* **66** (1991) 1126.

[44] M. Alcubierre, *The warp drive: hyper-fast travel within general relativity*, *Class. Quant. Grav.* **11** (1994) L73.

[45] K. Gödel, *An example of a new type of cosmological solutions of Einstein's field equations of gravitation*, *Rev. Mod. Phys.* **21** (1949) 447.

[46] M.M. Som and A.K. Raychaudhuri, *Cylindrically Symmetric Charged Dust Distributions in Rigid Rotation in General Relativity*, *Proc. Roy. Soc. London A* **304** (1968) 81.

[47] A.K. Raychaudhuri and S.N. Guha Thakurta, *Homogeneous space-times of the Gödel-type*, *Phys. Rev. D* **22** (1980) 802.

[48] M.J. Rebouças, J.E. Åman and A.F.F. Teixeira, *A note on Gödel-type space-times*, *J. Math. Phys.* **27** (1986) 1370.

[49] M.J. Rebouças and A.F.F. Teixeira, *Features of a Relativistic Space-Time with Seven Isometries*, *Phys. Rev. D* **34** (1986) 2985.

[50] F.M. Paiva, M.J. Rebouças and A.F.F. Teixeira, *Time Travel in the Homogeneous Som-Raychaudhuri Universe*, *Phys. Lett. A* **126** (1987) 168.

[51] A. Krasiński, *Rotating Dust Solutions of Einsteins Equations With 3-Dimensinal Symmetry Groups - III - All Killing Fields Linearly Independent of U(Alpha) And W(Alpha)*, *J. Math. Phys.* **39** (1998) 2148.

[52] M. Tsamparlis, D. Nikolopoulos and P.S. Apostolopoulos, *Computation of the conformal algebra of 1 + 3 decomposable spacetimes*, *Class. Quantum Grav.* **15** (1998) 2909.

[53] S. Carneiro, *A Gödel-Friedman cosmology?*, *Phys. Rev. D* **61** (2000) 083506.

[54] Y.N. Obukhov, *On physical foundations and observational effects of cosmic rotation*, in 'Colloquium on Cosmic Rotation' Eds. M. Scherfner, T. Chrobok and M. Shefaat, pp 23–96 (Wissenschaft und Technik Verlag: Berlin, 2000) *astro-ph/0008106*.

[55] S. Carneiro and G.A. Mena Marugan, *Anisotropic cosmologies containing isotropic background radiation*, *Phys. Rev. D* **64** (2001) 083502.

[56] J.D. Barrow and C.G. Tsagas, *Dynamics and stability of the Gödel universe*, *Class. Quantum Grav.* **21** (2004) 1773.

[57] M.P. Dabrowski and J. Garecki, *Energy momentum and angular momentum of Gödel universes*, *Phys. Rev. D* **70** (2004) 043511.

[58] F.C. Sousa, J.B. Fonseca and C. Romero, *Equivalence of three-dimensional spacetimes*, *Class. Quantum Grav.* **25** (2008) 035007.

[59] L.L. Smalley, *Gödel cosmology in Riemann-Cartan spacetime with spin density*, *Phys. Rev. D* **32** (1985) 3124.

[60] J. Duarte de Oliveira, A.F.F. Teixeira and J. Tiomno, *Homogeneous Cosmos of Weyssenhoff Fluid in Einstein-Cartan Space*, *Phys. Rev. D* **34** (1986) 3661.

[61] A.J. Accioly and G.E.A. Matsas, *Are there causal vacuum solutions with the symmetries of the Gödel universe in higher-derivative gravity?*, *Phys. Rev. D* **38** (1988) 1083.

[62] J.D. Barrow and M.P. Dabrowski, *Gödel universes in string theory*, *Phys. Rev. D* **58** (1998) 103502.

[63] J.E. Åman, J.B. Fonseca-Neto, M.A.H. MacCallum and M.J. Rebouças, *Riemann-Cartan spacetimes of Gödel-type*, *Class. Quantum Grav.* **15** (1998) 1089.

[64] M.J. Rebouças and A.F.F. Teixeira, *Riemannian Space-times of Gödel Type in Five Dimensions*, *J. Math. Phys.* **39** (1998) 2180.

[65] M. J. Rebouças and A.F.F. Teixeira, *Causal Anomalies in Kaluza-Klein Gravity Theories*, *Int. J. Mod. Phys. A* **13** (1998) 3181.

[66] P. Kanti and C.E. Vayonakis, *Gödel-type universes in string-inspired charged gravity*, *Phys. Rev. D* **60** (1999) 103519.

[67] H.L. Carrion, M.J. Rebouças and A.F.F. Teixeira, *Gödel-type Spacetimes in Induced Matter Gravity Theory*, *J. Math. Phys.* **40** (1999) 4011.

[68] E.K. Boyda, S. Ganguli, P. Horava and U. Varadarajan, *Holographic protection of chronology in universes of the Gödel type*, *Phys. Rev. D* **67** (2003) 106003.

[69] J.D. Barrow and C.G. Tsagas, *The Gödel brane*, *Phys. Rev. D* **69** (2004) 064007.

[70] M. Banados, G. Barnich, G. Compere and A. Gomberoff, *Three-dimensional origin of Gödel spacetimes and black holes*, *Phys. Rev. D* **73** (2006) 044006.

[71] W.-H. Huang, *Instability of Tachyon Supertube in Type IIA Gödel Spacetime*, *Phys. Lett. B* **615** (2005) 266.

[72] D. Astefanesei, R.B. Mann and E. Radu, *Nut Charged Space-times and Closed Timelike Curves on the Boundary*, *JHEP* **01** (2005) 049.

[73] Y. Brihaye, J. Kunz and E. Radu, *From black strings to black holes: nuttier and squashed AdS_5 solutions*, *JHEP* **08** (2009) 025.

[74] C. Furtado et al., *Gödel solution in modified gravity*, *Phys. Rev. D* **79** (2009) 124039.

[75] M.J. Rebouças and J. Santos, *Gödel-type universes in f(R) gravity*, *Phys. Rev. D* **80** (2009) 063009.

[76] J. Santos, M.J. Rebouças and T.B.R.F. Oliveira, *Gödel-type universes in Palatini f(R) gravity*, *Phys. Rev. D* **81** (2010) 123017.

[77] A.A. Sousa, R.B. Pereira and A.C. Silva, *Energy and angular momentum densities in a Gödel-type universe in teleparallel geometry*, *Grav. Cosmol.* **16** (2010) 25.

[78] C. Furtado, J.R. Nascimento, A. Yu. Petrov and A.F. Santos, *Horava-Lifshitz gravity and Gödel Universe*, *Phys. Rev. D* **84** (2011) 047702.

[79] Z. Tao, W. Pu-Xun and Yu Hong-Wei, *Gödel-type universes in f(R) gravity with an arbitrary coupling between matter and geometry*, *Chin. Phys. Lett.* **28** (2011) 120401.

[80] C. Furtado et al., *The Gödel metric in the Chern-Simons modified gravity*, *Int. J. Mod. Phys. Conf. Ser.* **18** (2012) 145.

[81] D. Liu, P. Wu and H. Yu, *Gödel-type universes in f(T) gravity*, *Int. J. Mod. Phys. D* **21** (2012) 1250074.

[82] J.B. Fonseca-Neto, A.Yu. Petrov and M.J. Rebouças, *Gödel-type universes and chronology protection in Hořava-Lifshitz gravity*, *Phys. Lett. B* **725** (2013) 412.

[83] J.R. Nascimento, A. Yu. Petrov, A.F. Santos and W.D.R. Jesus, *Gödel solution in the bumblebee gravity*, *Mod. Phys. Lett. A* **30** (2015) 1550011.

[84] A.F. Santos and C.J. Ferst, *Gödel-type solution in f(R,T) modified gravity*, *Mod. Phys. Lett. A* **30** (2015) 1550214.

[85] P.J. Porfírio et al., *Chern-Simons modified gravity and closed timelike curves*, *Phys. Rev. D* **94** (2016) 044044.

[86] J.A. Agudelo et al., *Gödel and Gödel-type universes in Brans-Dicke theory*, *Phys. Lett. B* **762** (2016) 96.

[87] M. Gürses and Ç. Şentürk, *Gödel-type metrics in Einstein-Aether theory II: nonflat background in arbitrary dimensions*, *Gen. Rel. Grav.* **48** (2016) 63.

[88] T. Azizi and E. Yaraie, *Gödel-type universes in Palatini $f(R)$ gravity with a non-minimal curvature-matter coupling*, *Int. J. Theor. Phys.* **55** (2016) 176.

[89] T. Clifton and J.D. Barrow, *The existence of Gödel, Einstein, and de Sitter universes*, *Phys. Rev. D* **72** (2005) 123003.

[90] M.J. Rebouças and J. Tiomno, *Homogeneity of Riemannian space-times of Gödel type*, *Phys. Rev. D* **28** (1983) 1251.

[91] F. Bampi and C. Zordan, *A note on Gödel's metric*, *Gen. Rel. Grav.* **9** (1978) 393.

[92] M.J. Rebouças and J.E. Åman, *Computer-aided study of a class of Riemannian space-times*, *J. Math. Phys.* **28** (1987) 888.

[93] A.F.F. Teixeira, M.J. Rebouças and J.E. Åman, *Isometries of homogeneous Gödel-type spacetimes*, *Phys. Rev. D* **32** (1985) 3309.

[94] L. Pogosian and A. Silvestri, *Pattern of growth in viable $f(R)$ cosmologies*, *Phys. Rev. D* **77** (2008) 023503.

[95] S.W. Hawking, *Chronology protection conjecture*, *Phys. Rev. D* **46** (1992) 603.